

IN THE CLAIMS:

The status of each claim that has been introduced in the above-referenced application is identified in the ensuing listing of the claims. This listing of the claims replaces all previously submitted claims listings.

1. (Original) A method for eliciting a T-cell mediated immune response in an animal, comprising administering to the animal a quantity of a composition including an extract of an egg obtained from a nonmammalian source animal, said extract comprising transfer factor generated by said source animal in a T-cell mediated immune response to at least one antigenic agent.
2. (Original) The method of claim 1, wherein said administering comprises administering to the animal a quantity of said composition with said extract comprising transfer factor molecules having molecular weights of about 4,000 Da to about 5,000 Da.
3. (Original) The method of claim 1, wherein said administering is effected orally.
4. (Original) The method of claim 1, wherein said administering is effected nasally.
5. (Original) The method of claim 1, wherein said administering is effected parenterally.
6. (Original) The method of claim 1, wherein said administering is effected topically.
7. (Currently amended) The method of claim 1, wherein said administering comprises preventing administering a sufficient quantity of said composition to cause an immune system of the animal from exhibiting a disease state associated with to elicit an immune response against an infection by a pathogen associated with said antigenic agent.

8. (Original) The method of claim 7, wherein said administering is effected before the animal is exposed to said pathogen.

9. (Original) The method of claim 7, wherein said administering is effected after the animal has been exposed to said pathogen.

10. (Original) The method of claim 7, wherein said administering also comprises administering to the animal said composition with said transfer factor comprising transfer factor molecules specific for at least one antigen of said pathogen.

11. (Currently amended) The method of claim 1, wherein said administering comprises ~~treating-administering~~ a sufficient quantity of said composition to treat a disease state associated with infection by a pathogen associated with said antigenic agent.

12. (Original) The method of claim 11, wherein said administering also comprises administering to the animal said composition with said transfer factor comprising transfer factor molecules specific for at least one antigen of said pathogen.

B2
13. (Previously presented) The method of claim 1, wherein said administering comprises administering to the animal said composition with said transfer factor comprising transfer factor molecules specific for at least one antigen of at least one antigenic agent.

14. (Original) The method of claim 1, wherein said administering comprises administering to the animal said composition with said transfer factor comprising transfer factor molecules specific for at least one antigen of at least one of Newcastle Virus, rubeola virus, mumps virus, rubella virus, Epstein-Barr Virus, hepatitis B virus, and *H. pylori*.

15. (Original) The method of claim 1, wherein said administering comprises administering said composition to a mammal.

16. (Original) The method of claim 1, wherein said administering comprises administering to the animal said composition with said egg extract comprising an extract of an avian egg.

17. (Original) The method of claim 1, wherein said administering comprises administering to the animal said composition with said egg extract comprising an extract of a non-avian egg.

18. (Original) The method of claim 1, wherein said administering comprises administering to the animal said composition with said egg extract comprising non-mammalian transfer factor.

19. (Original) The method of claim 1, wherein, following said administering, said transfer factor causes the animal, *in vivo*, to elicit the T-cell mediated immune response.

B2
20. (Original) A method for causing an animal to elicit a T-cell mediated immune response, comprising:
administering to the animal a quantity of a composition including an extract of an egg obtained from a nonmammalian source animal, said extract comprising transfer factor generated by said source animal in a T-cell mediated immune response to at least one antigenic agent;
and
permitting the transfer factor and the animal's immune system to initiate the T-cell mediated immune response *in vivo*.

21. (Original) The method of claim 20, wherein said administering comprises administering to the animal a quantity of said composition with said extract comprising transfer factor molecules having molecular weights of about 4,000 Da to about 5,000 Da.

22. (New) The method of claim 1, wherein said administering comprises administering to the animal a sufficient quantity of said composition to strengthen an immune system of the animal.

B2
